



TITLE:

A small remark on finite multiple zeta values and p -adic multiple zeta values (Various aspects of multiple zeta values)

AUTHOR(S):

Yamashita, Go

CITATION:

Yamashita, Go. A small remark on finite multiple zeta values and p -adic multiple zeta values (Various aspects of multiple zeta values). 数理解析研究所講究録別冊 2017, B68: 171-174

ISSUE DATE:

2017-10

URL:

<http://hdl.handle.net/2433/243724>

RIGHT:

© 2017 by the Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

A small remark on finite multiple zeta values and p -adic multiple zeta values

By

Go YAMASHITA*

Abstract

We show a relation between finite multiple zeta values of depth 2 and p -adic multiple zeta values of depth 1.

Recently, Kaneko-Zagier defined finite multiple zeta values (i.e., a finite analogue of multiple zeta values), and they studied a (possible) relationship between the finite multiple zeta values and the (classical) multiple zeta values. In this short paper, we study a relationship between the finite multiple zeta values and the p -adic multiple zeta values in a little bit. More concretely, we show a relation between finite multiple zeta values of depth 2 and p -adic multiple zeta values of depth 1. This is a small partial answer to the question which the author was privately asked by Francis Brown in 2012 and by Masanobu Kaneko in 2013 independently (the question is on the relationship between finite multiple zeta values and p -adic multiple zeta values). The result comes from just a combination of calculations of Hoffman ([H]) and calculations in Washington's book ([W1997]), however, it seems to the author that no one has pointed out such relation yet and that no concrete relation between finite multiple zeta values and p -adic multiple zeta values is found yet within his best knowledge.

For $k_1, \dots, k_d \in \mathbb{Z}$, we define *mod p multiple zeta value* $\zeta_{\text{mod } p}(k_1, \dots, k_d) \in \mathbb{F}_p$ to be

$$\zeta_{\text{mod } p}(k_1, \dots, k_d) := \sum_{0 < n_1 < \dots < n_d < p} \frac{1}{n_1^{k_1} \dots n_d^{k_d}} \pmod{p}.$$

Received 24 February 2014. Revised June 29, 2014

2010 Mathematics Subject Classification(s): 11M38, 11S80

Key Words: finite multiple zeta values, p -adic multiple zeta values, Bernoulli numbers, Kummer congruence.

Supported by TOYOTA Central R&D Labs., Inc.

*RIMS, Kyoto University, Kyoto 606-8502, Japan.

e-mail: gokun@kurims.kyoto-u.ac.jp

Put $\mathcal{A} := (\prod_{p:\text{prime}} \mathbb{F}_p) / (\oplus_{p:\text{prime}} \mathbb{F}_p)$. Then, Kaneko-Zagier defined *finite multiple zeta value* $\zeta_{\mathcal{A}}(k_1, \dots, k_d) \in \mathcal{A}$ to be

$$\zeta_{\mathcal{A}}(k_1, \dots, k_d) := (\zeta_{\text{mod } p}(k_1, \dots, k_d))_{p:\text{prime}} \in \mathcal{A}.$$

On the other hand, Furusho defined *p-adic multiple zeta value* $\zeta_p(k_1, \dots, k_d)$ for $k_1, \dots, k_{d-1} \geq 1$, and $k_d > 1$ by using Coleman's *p*-adic integration theory (See [F2004], [Y2012] for the definition, [Y2010] for a fundamental conjecture and related theorem). (He also partially defined in the case where $k_d = 1$. However, we do not use this fact in this paper.)

We will show the following small remark:

Proposition 0.1. *Let $p > 2$ be a prime number, $n > 1$ a natural number such that $p-1$ does not divide $n-1$. Let B_m denote the m -th Bernoulli number (or Bernoulli-Seki number) for $m \in \mathbb{Z}_{\geq 0}$, i.e., the rational number defined by $\frac{t}{e^t-1} = \sum_{n \geq 0} B_n \frac{t^n}{n!}$. Set ord to be the p -adic valuation normalised by $\text{ord}(p) = 1$. Then we have*

1. $\text{ord}(\zeta_p(n)) \geq n$, and
2. For any $i > 0$ such that $p-1$ does not divide $i-1$, we have

$$\frac{1}{p^n} \zeta_p(n) \equiv \frac{(-1)^{n-i+1}}{\binom{n'}{i'}} \zeta_{\text{mod } p}(i, n-i) \equiv \frac{B_m}{m} \pmod{p},$$

where $m > 1$ is a natural number such that $m \equiv 1-n \pmod{p-1}$ (e.g. $m = p-n$ if $n < p-1$), and i', n' are natural numbers such that $0 < i', n' < p-1$ and $i' \equiv i \pmod{p-1}$, $n' \equiv n \pmod{p-1}$.

Remark. By the Kummer congruence, $\frac{B_m}{m}$ modulo p is independent of the choice of m .

Corollary 0.2. *For $n \in \mathbb{Z}_{>1}$ and $0 < i < n$, we have*

$$\zeta_{\mathcal{A}}(i, n-i) = \left[(*)_{p \leq \max(2, n+1)}, \left((-1)^{n-i+1} \frac{\binom{n}{i}}{p^n} \zeta_p(n) \pmod{p} \right)_{p > \max(2, n+1)} \right] \text{ in } \mathcal{A},$$

where $*$ is any element in \mathbb{F}_p , and $[]$ means modulo $\oplus_p \mathbb{F}_p$ (note that the class does not depend the choices of $*$ in \mathbb{F}_p for $p \leq \max(2, n+1)$ (i.e., finitely many p)). In particular, if $\zeta_{\mathcal{A}}(i, n-i)$ does not vanish, then $\zeta_p(n)$ does not vanish for infinitely many p .

Remark. For odd n , the non-vanishing of $\zeta_p(n)$ is equivalent to the higher Leopoldt conjecture (See [F2004, Examples 2.19 (b)]), which is currently known in the case where p is regular or $p-1$ divides $n-1$. Note also that it is not known yet that there is infinitely many regular primes.

Remark. For fixed $k \in \mathbb{Z}$, if $p - 1$ does not divide k , then we have $\zeta_{\text{mod } p}(k) = 0$. Therefore, we have $\zeta_{\mathcal{A}}(k) = 0$ for any $k \in \mathbb{Z}$. It is a general expectation of Kaneko-Zagier that the finite multiple zeta values of depth $d + 1$ behave like the multiple zeta values of depth d . The above proposition and corollary show a relation with the p -adic ones for $d = 1$.

Remark. Recently [IKT2014, p.11] shows that $\zeta_{\mathcal{A}}(1, \dots (n - 2 \text{ times}) \dots, 1, 2) = (B_{p-n} \bmod p)_p$. By this result, we also deduce that if $\zeta_{\mathcal{A}}(1, \dots (n - 2 \text{ times}) \dots, 1, 2)$ does not vanish, then $\zeta_p(n)$ does not vanish for infinitely many p .

Proof of the proposition. Take i' and n' as in the proposition. Then we have $\zeta_{\text{mod } p}(i, n - i) = \zeta_{\text{mod } p}(i', n' - i')$. We also have (see [H, Theorem 6.1])

$$\zeta_{\text{mod } p}(i', n' - i') = (-1)^{n' - i'} \binom{n'}{i'} \frac{B_{p-n'}}{n'} = (-1)^{n-i+1} \binom{n'}{i'} \frac{B_{p-n'}}{p - n'}$$

in \mathbb{F}_p , by using the formula of Bernoulli-Seki ([S1712], [B1713])

$$\sum_{0 < a < n} a^k = \frac{1}{k+1} \sum_{j=0}^k \binom{k+1}{j} B_j n^{k+1-j}.$$

Let $\omega : \mathbb{F}_p^\times \rightarrow \mathbb{Z}_p^\times$ be the Teichmüller character. Then, we have

$$\zeta_p(n) = \frac{p^n}{p^n - 1} L_p(n, \omega^{1-n})$$

for $n > 1$ by [F2004, Examples 2.19]. On the other hand, we have

$$L_p(a, \omega^{1-n}) \in \mathbb{Z}_p$$

for $a, n \in \mathbb{Z}$ such that $n - 1 \not\equiv 0 \pmod{p-1}$, by [W1997, Theorem 5.12], and we have

$$L_p(a, \omega^{1-n}) \equiv L_p(b, \omega^{1-n}) \pmod{p}$$

for $a, b, n \in \mathbb{Z}$ such that $n - 1 \not\equiv 0 \pmod{p-1}$, by [W1997, Corollary 5.13]. Thus, we have

$$\text{ord}(\zeta_p(n)) = n + \text{ord}(L_p(n, \omega^{1-n})) \geq n,$$

and

$$\frac{1}{p^n} \zeta_p(n) \equiv -L_p(n, \omega^{1-n}) = -L_p(n, \omega^m) \equiv -L_p(1-m, \omega^m) = (1-p^{m-1}) \frac{B_m}{m} \equiv \frac{B_m}{m} \pmod{p}$$

for $m > 1$ such that $m \equiv 1 - n \pmod{p-1}$. \square

Remark. By a repeated use of the formula of Bernoulli-Seki, we can show that $\zeta_{\text{mod } p}(k_1, \dots, k_{d+1})$ is a linear combination of products of d Bernoulli numbers by the same argument of the calculation of $\zeta_{\text{mod } p}(i, k-i)$ in [H].

We conjecture that $\text{ord}(\zeta_p(k_1, \dots, k_d)) \geq k_1 + \dots + k_d$ and $\frac{1}{p^{k_1 + \dots + k_d}} \zeta_p(k_1, \dots, k_d) \pmod p$ should be related to the finite multiple zeta values of depth $d+1$. The author was informed by Seidai Yasuda that Kentaro Ihara and he had formulated a precise conjectural relation between the finite multiple zeta values and the p -adic multiple zeta values for arbitrary depth.

Acknowledgement The author heartily thanks the referee for kindly informing him the reference of [H] and [IKT2014]. He thanks *Masanobu Kaneko* for introducing him to the theory of the finite multiple zeta values. He also thanks *Francis Brown* and *Masanobu Kaneko* for asking me the question on the relation between the finite multiple zeta values and the p -adic multiple zeta values. He sincerely thanks *TOYOTA Central R&D Labs. Inc.* for offering him a special position in which he can concentrate on pure math research. He also sincerely thanks *Sakichi Toyoda* for his philosophy, and the executives of TOYOTA Central R&D Labs. Inc. for inheriting it from him for 80 years after his death.

References

- [B1713] Bernoulli, Jakob, *Ars Conjectandi, opus posthumum. Accedit Tractatus de seriebus infinitis, et epistola gallicé scripta de ludo pilae reticularis*. Basel, Thurneysen Brothers, 1713.
- [F2004] Furusho, Hidekazu, *p -adic multiple zeta values. I. p -adic multiple polylogarithms and the p -adic KZ equation*. Invent. Math. **155** (2004), no. 2, 253–286.
- [IKT2014] Imatomi, Kohtaro; Kaneko, Masanobu; Takeda, Erika, *Multi-Poly-Bernoulli Numbers and Finite Multiple Zeta Values*. Journal of Integer Sequences. **17**, 1–12, (2014)
- [H] Hoffman, Michael E. *Quasi-symmetric functions and mod p multiple harmonic sums*. preprint, <http://arxiv.org/pdf/math/0401319v2.pdf>
- [S1712] Seki, Kōwa, *Katsuyōsampo*. Edo(Tokyo), Masuya Gorōwemon, 1712.
- [W1997] Washington, Laurence, *Introduction to Cyclotomic Fields*. Springer-Verlag, New York, 1997.
- [Y2010] Yamashita, Go, *Bounds for the dimensions of p -adic multiple L -value spaces*. Documenta Math. Extra Volume: Andrei A. Suslin's Sixtieth Birthday (2010), 687–723.
- [Y2012] Yamashita, Go, *p -adic multiple zeta values, p -adic multiple L -values, and motivic Galois groups*. Adv. Studies in Pure Math. **63** (2012), 629–658.